

Brief Documentation for Boxwood Blight Infection Risk Model at Uspest.org

Version 1.1 - July 6, 2016

Leonard Coop, coopl@science.oregonstate.edu
Associate Director for Decision Support Systems
Integrated Plant Protection Center
Oregon State University
<http://uspest.org/wea>

Once reviewed and sufficiently verified, the model will be officially released and added to the multi-pest modeling website as per the link:

http://uspest.org/risk/models?mdl=bxwd_s

(note bxwd_s is the key standing for “boxwood blight – susceptible varieties”)

This is a summary of key modeling information developed within the spreadsheet: model_boxwood_blight_v11.xls and PDF version: model_boxwood_blight_v11.pdf

Abstract

The recently introduced pathogen, *Cylindrocladium pseudonaviculatum* (also *C. buxicola*) causes boxwood blight on boxwood, *Buxus sempervirens*. Models to predict infection and severity of this disease are urgently needed. As part of the IPPC/USPEST.ORG commitment to IPM and tools to mitigate the impact of invasive pests in the US, an infection risk model using the “degree-hours (DH) during periods of leaf wetness” approach was developed in order to better predict environmental conditions conducive to infection. This model was developed from recently presented data, thus far from two main sources; 1) a presentation by Belgian researchers Gehesquiere, Huylensbroeck, and Heungens (Sept. 2012), for two varieties of *Buxus*; *B. sempervirens* (Common or American boxwood), which has been classified as susceptible, and *B. sempervirens* var. *Suffruticosa* (English or true dwarf boxwood), which was found to be among the most highly susceptible varieties assessed (see http://go.ncsu.edu/boxwood_blight_links); and 2) boxwood blight webinars available (Mar. 2012 & Mar. 2013) from <http://www.anla.org>, including presentations by Sharon Douglas (Conn. Agric. Exp. Sta.) and others. The model will be revised as additional data become available. Two lower temperature thresholds were fit using the x-intercept regression method: 46F (7.78C), for first infections on young foliage; and 51F (10.56C), for first infections on mature foliage. First infection DH values (in Fahrenheit) were: Young leaves (T_{low}=46F): *B. s. var. Suff.*: 56 DHs, *B. s.*: 160 DHs; Mature leaves (T_{low}=51F): *B. s. var. Suff.*: 41 DHs, *B. s.*: 144 DHs. Since our goal for this infection risk model is to provide early warning of first infections of most sensitive plants (in this case to young foliage), the 46F/7.78C threshold was used for the single model developed and implemented at the website thus far. The model was also extended to include the relative degree of infection (measured as the number of lesions per plant reported by Gehesquiere et al.). Linear regression models were fitted to these data (young plus mature foliage combined) using the 7.78C threshold. Selected additional predictions (in Fahrenheit) of this extended model include: at 250 DHs: *B. s. var. Suff.*: 6 lesions, *B. s.*: 1 lesion; at 400 DHs: *B. s. var. Suff.*: 12 lesions, *B. s.*: 3 lesions; at 550 DHs: *B. s. var. Suff.*: 18 lesions, *B. s.*: 5 lesions. The model was implemented as version 1.0 and later updated as version 1.1 at http://uspest.org/risk/models?mdl=bxwd_s. Early tests indicate that DH values accumulate in a pattern similar to apple scab, but that box blight infection sensitivity is overall much lower than apple scab during warm/wet periods (Figs 4a and 4b). The model may need further updates as additional results are released, including covering a wider range of susceptibilities beyond the two boxwood varieties studied thus far.

Models of First Infection

Time of first symptoms of infection (lesions on leaves), (data from Gehesquiere et al. 2012, Fig. 1) were plotted vs. temperature (C) to determine potential lower thresholds, both for infection of young leaves and mature (less susceptible) leaves, for the two varieties of *Buxus* studied, *B. sempervirens* and *B. s. var. Suffruticosa* (Fig. 2).




Fig. 1. Data of Gehesquiere et al. 2012 used to develop *Buxus* infection risk models.




Fig. 2. Regression of rate of first lesion development (1/hours) vs. Temperature (C) for two varieties of boxwood, and for young leaves (Top), and mature leaves (Bottom).

These models indicated low temperature thresholds (using the x-intercept method) for first lesions of between 7.0 and 8.1 C for young leaves, and 12.4 and 12.8 C for mature leaves for the two varieties of boxwood. When combined, and forced to the nearest whole integer (in F) and extended past initial infections (see below), these models were modified to a low threshold of 7.78 C (46 F) for young leaves, and to 10.56 C (51 F) for mature leaves. The modified regression equations retained a reasonably good fit to the data for first infection models ($r^2 = 0.92$ and 0.79 , young leaves; 0.93 and 0.97 , mature leaves). The resulting DHs to first lesions were: Young leaves ($T_{low}=46$ F): B. s. var. Suff.: 56 DHs, B. s.: 160 DHs; Mature leaves ($T_{low}=51$ F): B. s. var. Suff.: 41 DHs, B. s.: 144 DHs.

Degree of Infection Models

From the data of Gehesquiere et al. (2012), we regressed the number of leaf lesions per plant (reflecting the number of infections) vs. DH during leaf wetness periods. Varying the low temperature threshold (T_{low}) for DH calculation and recording r^2 values to indicate best fit indicated that a T_{low} in the range of 10.5-11 C was best for the data when the young leaves, shown above to be more susceptible at lower temperatures, were included in the analysis, and 11.5-12 C when young leaves were excluded. This degree-of-fit analysis was repeated after dropping the 3 data points with highest number of lesions (due to a leveling off/non-linear effect). Highest r^2 values were obtained for T_{lows} of 11-11.5 C when including young leaves, and 12 C excluding young leaves.

While these higher T_{low} values provide a better fit when infection of mature leaves is modeled, we set an overall common threshold for all models to the lower value of 7.78 C (46 F) to meet the following objectives: 1) Develop a single (pair of) conservative models that best predict first infections on young, more susceptible leaves (for both varieties), 2) Include additional degree of infection risk estimates for young plus mature leaves, 3) Working with the data revealed that the experimental conditions went far beyond the leaf wetness/degree-hour duration expected under natural outdoor conditions, so the highest levels of infection (>500 DHs) were dropped from the analysis, 4) Provide a whole number T_{low} when converted to degrees F to facilitate implementation of the model. Using the 7.78 C low threshold, two models of disease build up were then fit to the data (Fig. 3):

B. semp.: Degree of infection (#lesions/plant) = $0.0243 \times DH - 2.2275$; $r^2 = 0.946$

B. s. v. Suff.: Degree of infection (#lesions/plant) = $0.0706 \times DH - 4.0446$; $r^2 = 0.877$




Fig. 3. Degree of infection models (converted to Fahrenheit) and data out to 600 DH.

Model Implementation at USPEST.ORG/WEA

The above models describing first infection risk and degree of infection have been combined and implemented for review and testing at the multi-pest modeling website http://uspest.org/risk/models?mdl=bxwd_s. It is, like the implementation of apple and pear scab models, a standard “degree-hours during periods of leaf wetness” model. The degree of infection model is subject to interpretation of events relevant to management needs, and therefore should be reviewed especially by potential end-users of the model. The combined model currently uses the following selected events and parameters drawn from the two degree of infection models:

Summary of Parameters for Modeling Boxwood Blight :

Name of model:	Boxwood blight infection risk
Model type:	Degree-hours (DHs) accum. during leaf wetness periods
Lower temp. threshold:	46F (7.78C)
Upper temp. threshold:	85F (29.4C)
No. of dry hours to stop the infection cycle:	more than 3.0 (reduced from 8.0 following recent research funded by USDA APHIS PPQ)
DHs to first infection of young leaves (highly susc. Var.):	56
DHs to first infection of young leaves (susc. Var.):	160
DHs for infection resulting in: 6 lesions, highly susc. Var., 1 lesion, susc. Var:	250
DHs for infection resulting in: 12 lesions, highly susc. Var., 3 lesions, susc. Var:	400
DHs for infection resulting in: 18 lesions, highly susc. Var., 5 lesions, susc. Var:	550
Model assumptions:	<ol style="list-style-type: none">1. Spores from microsclerotia generally require rainfall to spread and initiate the infection process, thus the model conservatively does not require rainfall events, as spores may also be present from existing lesions.2. The model should reflect a range of infection conditions most likely to occur in typical N. America climates; it was adjusted to reflect needs in the humid mid-latitudes (such as NC, VA, WV, PA, and MD).3. These results reflect work performed on one highly susceptible (English boxwood) and one susceptible (American boxwood) variety; lower infection risk levels would be expected for less susceptible varieties.

Initial tests of model parameters were compared to the apple scab model using the website at <http://uspest.org/risk/models>:




Fig. 4a. Comparison of preliminary infection risk model V1.1 online at http://uspest.org/risk/models?mdl=bxwd_s&mdl=a_scab, for boxwood blight (bottom plot) vs. apple scab infection risk model (middle plot), May 10-Jun 9, 2012, Blacksburg, VA. Note that the boxwood blight reset threshold was changed from 8 dry hours to 3 dry hours based on new research since version 1.0.




Fig. 4b. Same time interval model output for CRVO Corvallis, Oregon. Note that boxwood blight infection conditions are less favorable than for apple scab.




Fig. 4c. The same preliminary model example output (as in Fig. 4a) in tabular format (requires clicking on the plot for the table to pop-up, note scroll bar indicating only partial results are displayed), at http://uspest.org/risk/models?mdl=bxwd_s.

Susceptibility of Commercial Varieties to Box Blight (analysis based on final disease assessment)




Fig. 5. Information on relative susceptibility of varieties of boxwood to boxwood blight. From NCSU website, http://go.ncsu.edu/boxwood_blight_links. For reference in interpreting current model outputs.

Areas for Potential Model Improvement

This model (version 1.1) has been in use since 2013 and has been updated once thus far. Other versions of the model which include wind and rain were developed, but at this time evidence does not support their use. This updated model is much less conservative than version 1.0, now that the infection process is halted after only a 3 hr dry period, vs 8 hr in the first version. The model has been partly validated by comparing model predictions prior to observed outbreaks in several instances. In all cases, the model gave sufficient warning that infection could spread if inoculum is present. If you would like to contribute to validation, please send reports of outbreaks to: coopl@science.oregonstate.edu . Please include in your posting: 1) the varieties of boxwood affected, 2) the time period when symptoms were first observed/noted to rapidly expand, the location (nearest town would be sufficient), and 3) a description of the setting: park or other public space, nursery, private home, etc. We will use your info. To run the model just prior/during the observed outbreaks to check that the model is sufficiently sensitive to environmentally-conducive conditions.

Additional forthcoming data relating environmental conditions to infection risk are expected to improve the model in the near future. In addition, the several other varieties of *Buxus*, as well as other susceptible plant groups should be compared to these results and perhaps the model can be revised or split for several categories of susceptibility, such as moderately tolerant, moderately susceptible, susceptible, and highly susceptible (as shown in Fig. 5.).